

# **Innovation in mathematics**



**Jim H. Adams**

[www.jimhadams.org](http://www.jimhadams.org)

Copyright © 2014 Jim H. Adams

This work may be translated or copied in whole or in part without the permission of the author, Jim H. Adams, provided that he is acknowledged as the source of the text, which is asserted by him unless indicated otherwise. Use is allowed in connection with any information storage and retrieval system, electronic adaption, computer software, or by any methodology now known or hereafter developed.

---

Jim H. Adams is a researcher in the concepts of mathematics and their explicit representation. This work is a guide to some of these ideas, innovations or developments. It is also an encouragement to the aspiring mathematician to get involved in mathematical research at an early stage, and continually.

---

After a career in IT, Jim joined New Music Brighton as a composer and performer of his own works. He has been extensively involved in mathematical research.

The eBook *The climate and energy emergencies* was published in 2014, this eBook was also concluded in 2014 and *Superexponential algebra* will be released in 2015.

## Foreword

This eBook describes some of my own works that constitute innovations in mathematics, a description, helpful to the interested reader, of the creative process in mathematics, and finally a translation into English of an article by Eisenstein that I have found full of astounding insight.

From the desire to understand the mathematics of the twentieth century, the eBook arose from the study of the mathematics of the nineteenth century and before, and progressively a commentary and some modifications of it. To the best of my knowledge, these innovations were never implemented in the twentieth century.

*Creative mathematics – a viewpoint* ties the themes of the book together. It is a light-hearted look at ways of creative thinking and implementing them in practice.

*Discussion on ladder numbers and zero algebras* provides for the general reader innovations in real number theory: a proof of the inconsistency of the uncountable continuum hypothesis versus countable Cauchy sequences for reals. Implicitly, this inconsistency has been thought a possibility since the work of Gödel. In more detail, a set  $S$  is countable with respect to a set  $T$  if there exists at least one bijection between  $S$  and  $T$ , but there may be other maps between the same sets which are not bijections. Using this definition of relative countability to look at Cantor's diagonal argument, we also give an example which does not generate a diagonal counterexample. We demonstrate by an extension of the proof of the countability of  $\mathbb{Q}$  that the set  $\{\mathbb{N}, \mathbb{N}^2, \dots, \mathbb{N}^p, \dots\}$  is countable and we further show that superexponential operations, a generalisation of exponentiation, applied to the set of natural numbers  $\mathbb{N}$ , always generate sets isomorphic to  $\mathbb{N}$ , thus deconstructing the existence of a distinct power set. We introduce ladder numbers, which are not well-ordered, implying the failure of a standard density argument of  $\mathbb{Q}$  within them. We use methods of transfinite induction acting on the density of  $\mathbb{Q}$  in  $\mathbb{R}$  to demonstrate the inconsistency of currently axiomatised analysis, and provide an alternative. Many uses of ladder numbers are sketched.

*Polynomial equations I – duplicate roots* adds a nice twist to Galois theory – the sextic with duplicate roots is solvable by radicals, even though the quintic is not solvable directly by using Galois techniques. We indicate ways Galois theory may be generalised rigorously.

*Polynomial equations II – transcendental solutions* gives the general algorithm for solving polynomial equations of any degree by successive convergent approximations.

*Introduction to intricate and hyperintricate numbers I* develops the well-known idea that  $\sqrt{-1}$  can be represented by the matrix  $\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$ . Since complex numbers have two components, but  $2 \times 2$  matrices have four, we develop a four dimensional object – an intricate number – that is an extension of complex numbers. This is generalised to  $2^n \times 2^n$  matrices – hyperintricate numbers. Proofs using the algebra are given also in *Introduction to intricate and hyperintricate numbers II*, which takes this development technically further.

*Some simple proofs of general reciprocity*, after proving the quadratic reciprocity theorem, introduces a minor extension of the Euler totient theorem. It then uses the work of Eisenstein to develop ideas further, including for totient cubic and quartic reciprocity. Implications for the exponential algebra D1, mentioned throughout the book, which differs from the standard in that  $(e^{i0})^i = e^{i0}$ , are also indicated.

*An elementary investigation of the prime  $p = 4k - 1$  asymmetry theorem for quadratic residues I* investigates by elementary techniques the theorem known to hold by sophisticated transcendental techniques that for prime  $p = 4k - 1$  the number of quadratic residues in the interval  $[1, 2k - 1]$  is greater than in the interval  $[2k, 4k - 2]$ . *Part II* of the same work introduces the concept of *parabolas* to describe this phenomenon. These elementary method ideas are related to the known result on the absence of a tenth discriminant, and a proof by these means would constitute a proof of this theorem by elementary methods.

Lastly, we provide a translation from French into English of Eisenstein's *Applications of algebra to transcendental arithmetic*.

I have found that to develop my results I have increasingly been using the undergraduate text *Transform linear algebra* by Frank Uhlig, and both from there and elsewhere his insights. I would like to thank a number of people for their comments on parts of the book, which changed some aspects of it: Roger Fenn, Doly García, Tim Gibbs, Roger Goodwin, Ben Greenfield, Paul Hammond and Steven Miller, and to James Hirschfeld, who stressed the importance of completing a project; this has become a permanent intention in my writing. The decisions made on the contents are my own, as is the responsibility for any errors.

Jim H. Adams

Brighton & Hove

2014

## **Supplementary notes, May 2015, April 2017 and March 2018**

Changes to this eBook are now frozen from March 2018 as a record of my thinking up to 2014. These supplementary notes were mainly created in May 2015. From April 2017 the work has contained Part III of *An elementary investigation of the prime  $p = 4k - 1$  asymmetry theorem for quadratic residues* and from March 2018 a proof of the asymmetry theorem, with corresponding renumbering of the final chapter and additions to the index.


Where references are made to the work *Superexponential algebra* before it was completed, these references have been updated, and elsewhere. No other modifications have been made.

Intricate numbers are more usually given the name split-quaternions or coquaternions. They were introduced by James Cockle in 1849 under the latter name in the London-Edinburgh-Dublin Philosophical Magazine.

The argument used in *Discussion on ladder numbers and zero algebras* can be made more precise on page 02.19.

For finite or countably infinite sets  $S$ ,  $T$  or  $U$  we define  $\equiv$  by the existence of at least one bijection within the natural numbers  $\mathbb{N}$ . This is an equivalence relation, in that  $S \equiv S$ , if  $S \equiv T$  then  $T \equiv S$ , and if  $S \equiv T$  and  $T \equiv U$ , then  $S \equiv U$ , so this forms a partition between those sets belonging to the equivalence class, and those outside it. Then for sets  $S_n, T_n, n \in \mathbb{N}$ , if for each  $n$   $S_n \equiv T_n$ , then for the set of all  $S_n, \{S_{nn}\} \equiv \{T_{nn}\}$ , where this means if  $S_n \equiv T_n$ , then the bijection is maintained for  $S_{n+1} \equiv T_{n+1}$ , and the second subscript indicates a distinguished copy for  $S_n \neq \emptyset$ , defined inductively:  $S_{n1} = \{S_n\}, S_{n2} = \{\{S_n\}\}, S_{np+1} = \{S_{np}\}$ .

In this work we will adopt the argument of Cantor that the set  $\mathbb{Q}$  of rationals is countable. Define the Cartesian product of all natural number pairs  $\mathbb{N} \times \mathbb{N}$  as  $\mathbb{N}^2$ . Consider the rational numbers not in lowest terms given by the set  $\mathbb{Q} \equiv \{\{1/n\}, \{2/n\}, \{3/n\}, \dots\} \equiv$  the unordered distinguished copies  $\{\mathbb{N}_1, \mathbb{N}_2, \mathbb{N}_3, \dots\} \equiv \mathbb{N} \times \mathbb{N}$  (a set of ordered pairs) which by the Cantor argument given next is  $\equiv \mathbb{N}$ . The mapping from  $\mathbb{N}$  to  $\mathbb{Q}$  is given in the following diagram.



What is meant by the symbols  $\dots$  in the sets just given? This indicates that if the  $p$ th position is occupied, then a similar item exists at position  $p + 1$ , although we can remove  $\dots$  from the language, and use the properties of  $\mathbb{N}$  itself. Then by induction defined through the properties of  $\mathbb{N}$ , we have  $\mathbb{N} \equiv \mathbb{N}^p$ , for  $p$  a natural number, so that the set  $\mathbb{N} \equiv \{\mathbb{N}_1, \mathbb{N}_2^2, \dots, \mathbb{N}_p^p, \dots\} \equiv \{\mathbb{N}_1, \mathbb{N}_2 \times \mathbb{N}_2, \dots, \mathbb{N}_p \times \mathbb{N}_p \times \dots (p \text{ terms}), \dots\}$  contains by definition  $\mathbb{N}^{\mathbb{N}} \equiv \{\mathbb{N} \times \mathbb{N} \times \dots\}$ , in violation of the basic assumption of the continuum hypothesis in set theory, that  $\{0, 1\}^{\mathbb{N}}$  is uncountable.

The description of the determinate zero algebra in the same paper is given on an axiomatic basis in the author's eBook *Superexponential algebra*.

The paper in session 11 and also the paper *Polynomial equations I – duplicate roots* assumes Galois *solvability* theory holds, which is investigated further in the eBook *Superexponential algebra*, where a description based on ring automorphisms is shown not to be equivalent to inner group automorphisms. We are not challenging Galois *representation* theory, a separate and distinct topic in multiplicative number theory. End results of Galois solvability theory are obtained from its replacement in the theory of varieties in the case of 'killing central terms', on eliminating dependencies.

# Table of Contents

## 1. Creative mathematics – a viewpoint.

1. Creative mathematical research and the system.
  - 1.1. Supporting creative intentions.
  - 1.2. Emancipation and control.
  - 1.3. The sociology of group innovation.
2. The creative mathematician.
  - 2.1. What is mathematics?
  - 2.2. Style.
  - 2.3. Intuition in mathematics.
  - 2.4. Errors.
  - 2.5. Methods for creative mathematical thinking.

## 2. Discussion on ladder numbers and zero algebras.

1. Session one – prologue.

### Foundations.

2. Session two – ladder algebras and nonstandard analysis.
3. Session three – the transfer principle.
4. Session four – ladder number counting and constructability.
5. Session five – ordering and the axiom of choice.
6. Session six – Cantor diagonalisation.
7. Session seven – transfinite induction and analysis.
8. Session eight – ordinals and cardinals.

### Applications.

9. Session nine – limits and convergence.
10. Session ten – ladder number transcendence.
11. Session eleven – Galois theory and infinite ladder automorphisms.
12. Session twelve – ladder calculus.
13. Session thirteen – further mathematical applications.
14. Session fourteen – zero algebras.

## 3. Polynomial equations I – duplicate roots.

1. Prologue.
2. Conceptual history.
  - 2.1. Public discourse.
  - 2.2. Personal reflections.

3. Results originating from the study of dependent roots.
  - 3.1. Polynomials of degree  $\leq 6$  with roots  $(x + a)(x - a)$ .
  - 3.2. Polynomials of degree  $\leq 6$  with duplicate roots.
  - 3.3. A differential condition for the detection of duplicate roots.
  - 3.4. The quintic with the root  $(x + a)$  and an adjoined root  $(x + a)$  or  $(x - a)$ .
4. Results on leaped Jordan-Hölder series using multifunctions.
  - 4.1. The cubic via  $(x + A + B + C)(x + A + \omega B + \omega^2 C)(x + A + \omega^2 B + \omega C) = 0$ .
  - 4.2. The sextic as a leaped series  $S_6 \triangleright G_3 \triangleright G_2 \triangleright 1$ .
5. Simple problems.

## **4. Polynomial equations II – transcendental solutions.**

1. Introduction.
2. Transcendental solutions to polynomial equations.
  - 2.1. Kronecker's theorem.
  - 2.2. Conversion of polynomials to real roots and coefficients.
  - 2.3. Transcendental determination of root moduli for any complex polynomial.
  - 2.4. Transcendental solutions of complex polynomials in the general case.

## **5. Introduction to intricate and hyperintricate numbers I.**

1. Intricate numbers.
2. The intricate Euler relations.
3. Hyperintricate numbers.
4. Exterior, interior and relative coefficient algebras.
5. Factorisation of intricate numbers.
6. Symmetric, antisymmetric and upper triangular matrices.
7. Intricate products under non-commutation.
8. Representations of quaternions by hyperintricate numbers.
9. Roots of intricate basis elements.
10. Composites.

## **6. Introduction to intricate and hyperintricate numbers II.**

1. Intricate and hyperintricate numbers.
2. Deriving hyperintricate basis element coefficients from a matrix.
3. Extricate and duplicate numbers.
4. The J representation.
5. Standard division algebras and exoctonions.
6. Compression and expansion.

7. The hyperintricate trace, norm and layer algebra.
8. J-diffeomorphisms for non-fixed J.
9. J-abelian intricate powers.
10. J-abelian hyperintricate determinants and inverses.
11. The general hyperintricate inverse.
12. The matrix symmetriser classification.
13. The  $\mathcal{JAF}$  spectral D1 exponential algebra.

## **7. Some simple proofs on general reciprocity.**

1. Fermat's little theorem and quadratic reciprocity.
2. Euler's totient formula.
3. The Frobenius automorphism, Fermat's and Euler's theorems.
4. Totient quadratic reciprocity.
5. Quadratic Eisenstein representations.
6. Totient general reciprocity.
7. General Eisenstein representations.
8. The complex algebraic Fermat and Euler theorems.
9. The Euler theorem for superexponential congruence arithmetic.
10. The complex and intricate Fermat and Euler theorems for ladder numbers.
11. The complex and intricate exponential algebra D1.

## **8. An elementary investigation of the prime $p = 4k - 1$ asymmetry theorem for quadratic residues I.**

1. Introduction.
2. Preliminary remarks.
3. Detailed proofs leading to the disparity formula.
4. Disparity formulas and a condition for positive total disparity.
5. Ancillary remarks.
6. Some constraints on -1 disparities up to row T.
7. Trajectories.

## **9. An elementary investigation of the prime $p = 4k - 1$ asymmetry theorem for quadratic residues II.**

1. Synopsis of Part I, introduction to Part II and a plan of Part III.
2. Parabolas for rows.
3. Unstratified single fragments for rows.
4. Trajectories and trajectory parabolas.
5. Parabola and single fragment computations for the row region.
6. Multiple fragments in the row region.



## **10. An elementary investigation of the prime $p = 4k - 1$ asymmetry theorem for quadratic residues III.**

1. The class number and the tenth discriminant problem.
2. Further disparity expressions from Part I.
3. Relations with results for  $q = 4k' + 1$ .
4. Fragment and parabola computations for the trajectory region.
5. Even power residues for  $p = 4k - 1$ .
6. Positive total disparity from the arithmetic average of squares.
7. Composite numbers.

## **11. Applications of algebra to transcendental arithmetic, by G. Eisenstein.**

1. Quadratic residues.
2. Biquadratic residues.
3. Remarks.

## **12. Index.**