


Number, space and logic

Volume I

Superexponential structures



Jim H. Adams

www.jimhadams.com

Copyright © 2018 Jim H. Adams

This work may be translated or copied in whole or in part without the permission of the author, Jim H. Adams, provided that he is given as the source of the text, which is asserted by him unless indicated otherwise. It can be used in information storage and retrieval systems, in electronic adaption, computer software, or by any method now known or later developed.

The cover artwork is by Ian Dell.

Jim H. Adams is a researcher in the concepts of mathematics and their explicit representation. This work is a guide to some of these ideas, and an encouragement to the graduate to develop mathematical research productively and continually.

After a career in IT, Jim joined New Music Brighton as a composer and performer of his own works. He has been extensively involved in mathematical research.

The eBooks published in 2014 were *The climate and energy emergencies* and *Innovation in mathematics*. Volumes I, II and III of *Superexponential algebra* were completed in 2017, and *Number, space and logic* is a mathematical sequel to that work.

Keywords: l-adic numbers, elliptic curves, modular forms, complex multiplication, Heegner number, Riemann hypothesis, Goldbach conjecture.

Foreword

Number, space and logic combines a textbook at graduate level with a research project. It may be thought of as a commentary and development of the mathematics of the late 20th and the 21st centuries, in the same way as the eBook *Superexponential algebra* [Ad15] has been for the period in the 20th century before that covered in this work. As a practical guide to this, the eBook includes the mathematical machinery necessary for a proof of the general Riemann hypothesis and from it the weak Goldbach conjecture. There is an exposition of the theory of branched spaces and a Gentzen-type proof of the consistency of analysis.

In order to present our work in a coherent way, it has been necessary to proceed top down, from the abstract to the specific, but so that this generality can be understood we first give an account of the meanings of these abstractions. First we describe our approach to the finite and the infinite, then how our generalisations of space fit into the theory of trees and amalgams. Finally we show how superexponential structures can be used to encapsulate these ideas and extend them to an overarching theory of mathematics.

In the chapter on *the meaning of the finite and the infinite* we include a description of standard and significant results in finite and discrete number theory, including on p-adic numbers. We also discuss nonstandard results. Firstly we discuss extended sets of natural numbers, which we call transnatural numbers. The essential feature of these is that, unlike our results in nonstandard set theory, these numbers are not bijective to the natural numbers, instead they have a transfinite number of members, but otherwise they share the properties of natural numbers. For instance they contain transfinite prime numbers and can be used to define transfinite rational numbers. We also deduce the inconsistency of the uncountable continuum hypothesis compared with the countable rational numbers. We give an account of ladder algebra which incorporates a reformulation and extension of nonstandard analysis.

The chapter on *the meaning of branched spaces* introduces us to our extension of the domain of the Euler-Poincaré characteristic used in topology. In its familiar form our description appears more restricted than usual, although other formulations are equivalent to it. In its extended form it describes types of topology which have no analogue in standard topology. The model we introduce is one of branched spaces. We can put it this way: when a point is removed from the equivalent of an n-branched line, it divides into n pieces. An *explosion* is a set that contains an infinite number of branches. Indeed these are the topological analogues of other behaviour we will meet in exponential and superexponential number systems.

Concerning *the meaning of superexponentiation*, which is the topic of the last chapter of the eBook *Superexponential algebra* [Ad15], this idea is an extension of the repetitive process which generates multiplication from addition, and exponentiation from multiplication. So we generate an (n + 1)th superexponential operation by repetition of an nth superexponential operation. In that book we discussed nonstandard exponential and superexponential algebras, which we describe here by the statement that they are unbranched compared with standard such algebras. A real number we say has a *supernorm* defining its magnitude under an exponential or superexponential operation which is similarly real. Thus superexponentiation on numbers has two features that are interesting to investigate: supernorms and branching.

Having built up the meaning of these abstractions, from their axiomatic formulation we then provide a detailed exposition of these ideas in reverse order: of superexponential structures, trees and amalgams, and the finite and the infinite, and look at some of their consequences.

Volume I in more detail

Transfinite number theory is considered for the transnatural numbers \mathbb{M}_t , where the theory is an extension of finite arithmetic. We prove the inconsistency of the uncountable continuum hypothesis with respect to the countability of the rationals. Then we extend results on the ladder algebra of ordinal arithmetic to \mathbb{M}_t .

The author and the reader

A central objective of this work is to encourage the graduate to produce mathematical ideas, and also to give assistance by providing a minimally fussy exposition of the research into number theory of the last fifty years. We are seeking to foster insight so that the reader can pursue further investigations into the technical literature in a spirit of an understanding of its background.

Except for marketing, our inclination is that a proof is only the final stage in the presentation of a mathematical idea. Proofs are necessary, but not sufficient.

The objective of creative mathematics is the production of ideas. These need caring support, and part of this objective is not just to replicate results, but to reconstruct mathematics from new principles, to have the resilience to deconstruct what has gone before if it is needed, and to extend results by an analysis of the core ideas of the subject as it currently stands. Thus it becomes feasible to develop generalised new methods to tackle outstanding problems.

I am aware that the first language of the reader may not be English, so I have looked at the text and removed high literary style. Technical terms are I hope well explained, and examples are given.

I have provided no exercises to work through the text, the reason being that these are no more needed for the development of creative skills than the memorisation of words and standard literary works is necessary for an active participation in becoming an author. My hope is that the diligent reader will provide what is necessary for constructive work, and that it is no longer necessary to give direction.

The contents of this work are my own, as is the responsibility for any errors.

Jim H. Adams

Brighton & Hove

January 2017

Table of contents

Volume I

Superexponential structures

Foreword

Table of contents

Mathematical terms

Chapter I

The meaning of the finite and the infinite

- 1.1 Introduction
- 1.2 The Peano axioms for the transnatural numbers, \mathbb{M}_t
- 1.3 The Euclidean algorithm
- 1.4 Fermat's little theorem
- 1.5 Euler's totient formula
- 1.6 Transp-adic numbers
- 1.7 The inconsistency of the uncountable continuum hypothesis for $\mathbb{N}^{\mathbb{N}}$
- 1.8 Ordinal (ladder) arithmetic in \mathbb{M}_t

Chapter II

The meaning of branched spaces

- 2.1 Introduction
- 2.2 The Euler-Poincaré characteristic
- 2.3 The familiar square, cylinder, torus and cube
- 2.4 Branched spaces
- 2.5 Models for branched lines and areas
- 2.6 Deformation retracts and orientation
- 2.7 Branched handles, crosscaps and surgery of branched spaces
- 2.8 Explosion boundaries
- 2.9 Trees and amalgams
- 2.10 The general polynomial
- 2.11 The Gauss-Bonnet theorem
- 2.12 Embeddings and projections

Chapter III

The meaning of superexponentiation

- 3.1 Introduction
- 3.2 Dw superexponential structures
- 3.3 Supernorms and branching
- 3.4 The number Λ
- 3.5 Nonassociative superexponential representations
- 3.6 Superexponential substructures and singularities

Chapter IV
Complex multiplication

4.1 Introduction

Chapter V
Modular forms

5.1 Introduction

Chapter VI
Elliptic curves

6.1 Introduction

6.2 Elliptic curves

6.3 Rational points on derived elliptic curves

Chapter VII
Nonconformal analysis

7.1 Introduction

7.2 Fourier series and convolutions

7.3 The Maurer-Cartan equation

7.4 The intricate analytic Cauchy-Riemann equations

7.5 J-diffeomorphisms and non-fixed J

7.6 The intricate Cauchy-Riemann equations are nonconformal

7.7 The novanionic analytic Cauchy-Riemann equations

7.8 The nonconformal split representation

7.9 The complex Cauchy integral formula

7.10 The novanionic Cauchy integral formula

7.11 The nonconformal split Cauchy integral formula

7.12 Picard's theorems

Volume II
Trees and amalgams

Foreword
Table of contents
Mathematical terms

Chapter I
Trees and amalgams

- 1.1 Introduction
- 1.2 Trees

Chapter II
Branched spaces and explosions

- 2.1 Introduction
- 2.2 The superexponential model
- 2.3 Topological branching and explosions

Chapter III
Surgery

- 3.1 Introduction
- 3.2 The idea of surgery on branched spaces
- 3.3 Surgery on unbranched spaces

Chapter IV
Sequent calculus and colour logic

- 4.1 Introduction
- 4.2 Introduction to colour sets and other set theories
- 4.3 Supervarieties and colour logic
- 4.4 Classical sequent calculus and Gentzen normal form
- 4.5 A colour model for modal logics
- 4.6 Colour sequents

Chapter V
Homology

- 5.1 Introduction
- 5.2 Our view on K theory

Chapter VI
Cohomology

- 6.1 Introduction
- 6.2 Étale cohomology
- 6.3 Some comments on motivic cohomology

Chapter VII

Homotopy

7.1 Introduction

Volume III
The finite and the infinite

Foreword
Table of contents
Mathematical terms

Chapter I
Number theory

- 1.1 Introduction
- 1.2 Finite arithmetic
- 1.3 Ladder algebra and transcendence
- 1.4 l -adic numbers
- 1.5 l -adic Fourier series
- 1.6 Gauss, Ramanujan and Kloosterman sums
- 1.7 The Rogers-Ramanujan identities
- 1.8 Theta functions

Chapter II
Class field theory

- 2.1 Introduction

Chapter III
The consistency of analysis

- 3.1 Introduction
- 3.2 $V = L$
- 3.3 Forcing
- 3.4 Gentzen's consistency theorem for arithmetic
- 3.4 A Gentzen-type proof of the consistency of analysis

Chapter IV
Fermat's last theorem

- 4.1 Introduction
- 4.2 A proof of Fermat's last theorem by elementary methods

Chapter V
Global zeta functions

- 5.1 Introduction
- 5.2 Γ -functions
- 5.3 The basic problem
- 5.4 Some observations on the Riemann hypothesis
- 5.5 L -series

Chapter VI

Local zeta functions

- 6.1 Introduction
- 6.2 Background and history of the Weil conjectures
- 6.3 The statement of the Weil conjectures
- 6.4 The projective line
- 6.5 Projective space
- 6.6 Elliptic curves
- 6.7 Weil cohomology
- 6.8 Grothendieck's proof of three of the four conjectures
- 6.9 Deligne's first proof of the Riemann hypothesis conjecture
- 6.10 Deligne's second proof

Chapter VII

The Goldbach conjecture

- 7.1 Introduction
- 7.2 The Helfgott proof of the weak Goldbach theorem
- 7.3 The proof via the Riemann hypothesis

Chapter VIII

The Borcherds-Hajason construction

- 8.1 Introduction

References and Index

References

Index

Mathematical terms

The following terms, symbols, ideas and definitions are used in the text. The arrangement is by ideas rather than alphabetic. This may be scanned as a more technical alternative to the details of the contents in the Foreword, or as a further introduction to the contents which follow in the eBook.

1. Sets S, T (chapter III).

\emptyset (*the empty set*). The set with no members.

\odot (*the void set*). The set satisfying a false condition.

\in (*belongs to*). If x is a member of a set S then $x \in S$.

\subset (*properly included in*). If a set S is included in a set T and S does not equal T .

\subseteq (*included in*). Inclusion, when $S = T$ is possible.

$\mathcal{C}S_T$ (*complement of S in T*). Those x not belonging to S but that belong to T , and $S \subset T$.

\cup (*union of sets*). If x belongs to S or x belongs to T then x belongs to $S \cup T$.

\cap (*intersection of sets*). If $x \in S$ and $x \in T$ then $x \in S \cap T$.

2. \mathbb{N} is the set $\{1, 2, 3 \dots\}$ of *positive whole numbers*, otherwise called *natural numbers*. If this set contains the element 0, we denote it in this eBook by $\mathbb{N}_{\cup 0}$. If we wish to emphasise that it does not contain zero, we use $\mathbb{N}_{\neq 0}$.

\mathbb{Z} (from the German Zahl for number) is the set $\{\dots, -3, -2, -1, 0, 1, 2, \dots\}$ of negative, zero or positive *integers*.

\mathbb{M}_t is the set of *transnatural numbers* for index t , satisfying the rules for \mathbb{N} . $\mathbb{M}_1 = \mathbb{N}$. There are no surjections $\mathbb{M}_t \rightarrow \mathbb{M}_{t+1}$ or injections $\mathbb{M}_t \rightarrow \mathbb{M}' \rightarrow \mathbb{M}_{t+1}$ for distinct \mathbb{M}' .

\mathbb{Q} is the set of *rational numbers* m/n , where $m \in \mathbb{Z}$ and $n \in \mathbb{N}$, for example $1/2$.

\mathbb{A} is used in this eBook as the set of *algebraic numbers*, sums and differences of numbers of the form p^q , where $p, q \in \mathbb{Q}$, but p and q together are not both zero, for example $1 + 2^3\sqrt[5]{-1}$.

3. *Congruence arithmetic (mod n)*. Finite or ‘clock’ arithmetic where whole numbers come back to themselves, so its set is $\{0, 1, \dots (n - 1)\}$ and $n \neq 0$.

Prime number. A whole number which only when divided by 1 and itself gives a whole number. Example: 7.

Totient ($\phi(s)$). For a natural number s as a product of primes $p, q, \dots r$ to powers $j, k, \dots m$, if $s = (p^j)(q^k)\dots(r^m)$ then $\phi(s) = s \left[\left(1 - \frac{1}{p}\right) \left(1 - \frac{1}{q}\right) \dots \left(1 - \frac{1}{r}\right) \right]$. Example: $\phi(20) = 8$.

4. *Abelian*. Occurs for a set with a general operation $+$ (not necessarily addition as usually understood) when $a + b = b + a$ always.

Commutative. Abelian, but generally written for \times rather than $+$.

Associative. Satisfying $a + (b + c) = (a + b) + c$, or $a(bc) = (ab)c$, etc.

5. *Real (or Eudoxus) numbers*, \mathbb{U} . (chapter VII, and *Discussion on ladder numbers and zero algebras* in the eBook *Innovation in mathematics*, [Ad14]).

Complex numbers, \mathbb{C} . Numbers of the form $a + bi$, where $a, b \in \mathbb{U}$ and $i = \sqrt{-1}$.

Gaussian integers. Complex numbers where a and b above are integers.

\mathbb{F} is a *field* (chapter III). It contains axioms (rules) for addition and multiplication. Examples could be the real (or Eudoxus) numbers \mathbb{U} and complex numbers \mathbb{C} .

\mathbb{Y} is a *zero algebra* (chapter III). This is similar to a field except for the existence of multizeros.

Exponential algebra, (chapters XII and XIII). Contains axioms for exponentiation.

$\mathbb{D}w$ *exponential algebra*, (chapter XIII). A nonstandard exponential algebra.

Superexponentiation (chapter XV). An operation, of which the first three are addition, multiplication and exponentiation, where the n th is obtained by repeating the $(n - 1)$ th.

Superexponential algebra. Contains axioms connecting superexponential n th operations for various n .

6. Implies. (exercise, chapter XI). For statements A and B, A implies B is only false when A is true and B is false.

Sufficient. A is sufficient for B means A implies B.

Necessary. A is necessary for B means A is implied by B (the same as B implies A).

7. Function (chapter III). A set of pairs, $\{x, f(x)\}$, all x of which have a value $f(x)$.

Injection. A mapping from all the sets $\{a, b\}$ to $\{f(a), f(b)\}$, where $f(a) \neq f(b)$ if $a \neq b$.

Surjection. A mapping where every $f(x)$ in the set $\{f(x)\}$ has a value from an x .

Bijection. A mapping which is simultaneously injective and surjective.

8. Magma, M (chapter III). A set with one binary operation, with no other properties specified.

Polymagma (chapter XV). Maps a number of copies of a set to itself.

Group, G (chapter III). This satisfies the multiplicative axioms for a field, except that multiplication may be noncommutative: $ab \neq ba$.

Schur multiplier. A multiplicative group on identifying 1 and -1 as one element.

Subgroup. A set of elements in a group which satisfies within itself all the properties of the containing group.

Order of a group. The number of elements (or members) in a group.

Homomorphism of groups is a surjective map $h: G \rightarrow G'$ of groups with $h(ab) = h(a)h(b)$.

Isomorphism of groups. A bijective homomorphism.

Automorphism of a group is an isomorphism of a group to itself.

Inner automorphism of a group is an automorphism of the form $x \leftrightarrow a^{-1}xa$.

Outer automorphism of a group. An automorphism which is not inner.

Normal subgroup is invariant under all inner automorphisms of the containing group G .

Simple group has no normal subgroups other than itself and 1.

9. Ring, A. Satisfies the additive and multiplicative axioms of a field, except there is no general division and multiplication may be noncommutative. Example: matrices.

Unital ring. A ring with a multiplicative identity, 1. We assume rings are unital.

Automorphism of a ring. (chapter X). A bijective map of a ring A , $H: A \leftrightarrow A$, where $H(ab) = H(a)H(b)$ and $H(a + b) = H(a) + H(b)$.

10. Matrix (plural matrices). An array of numbers $B = b_{jk}$, where the element b_{ij} exists in the i^{th} row and j^{th} column. (chapters I and II).

Symmetric matrix. $U = u_{jk} = u_{kj}$.

Antisymmetric matrix. $V = v_{jk} = -v_{kj}$.

Matrix transpose. If $W = w_{jk}$, then the transpose $W^T = w_{kj}$.

Unit diagonal matrix. Denoted by $I = b_{jk}$, where $b_{jk} = 1$ when $j = k$, otherwise $b_{jk} = 0$.

Trace of a matrix. The sum of all (main) diagonal entries b_{jk} , where $j = k$.

Determinant (or hypervolume) of a matrix, ($\det B$). (chapters I and II).

Singular matrix, D. Satisfies $\det D = 0$.

11. Intricate number. A representation of 2×2 matrices, that is, with two rows and two columns, given by $a1 + bi + c\alpha + d\phi$. (chapter I).

Intricate basis element. One of the vectors $1, i, \alpha$ or ϕ above.

Real basis element. The number 1 in its intricate representation.

Imaginary basis element. The number i in its intricate representation.

Actual basis element. The number α in its intricate representation.

Phantom basis element. The number ϕ in its intricate representation.

Intricate conjugate. The number $a1 - bi - c\alpha - d\phi$.

J. $J = bi + c\alpha + d\phi$ in which $J^2 = 0$ or ± 1 .

\mathcal{JAF} . A changed basis for i, α and ϕ .

12. Hyperintricate number. A representation of $2^n \times 2^n$ matrices. (chapter II).

Layer. For example, a hyperintricate number with a component in 3 layers is $A_{B,C}$ where A, B and C are intricate numbers, possibly intricate basis elements.

n-hyperintricate number. A hyperintricate number representable by sums of components in n layers. Sometimes denoted by \mathfrak{Y}_n .

n-hyperintricate conjugate, \mathfrak{Y}_n^ .* Satisfies $\mathfrak{Y}_n^* \mathfrak{Y}_n = \det \mathfrak{Y}_n$.

J-abelian hyperintricate number. A number giving the example $A_B + \dots + D_E$, where $A = p1 + qJ, B = p'1 + q'J', \dots, D = t1 + uJ, E = t'1 + u'J'$. Two such hyperintricate numbers with identical J and J' commute.

Compression. The map $\kappa: A_B \rightarrow AB$.

Expansion. A map $\kappa^{op}: AB \rightarrow A_B$.

13. Vector, \mathbf{v} (in bold). A matrix as one row (a row vector), or as one column (a column vector). Example: the row vector (x, y, z) .

Scalar product of two vectors. The matrix product of multiplying each element of a row vector in turn with the corresponding elements of a column vector. Example: $x^2 + y^2 + z^2$.

Eigenvector. A vector \mathbf{x} satisfying for matrix $B, B\mathbf{x} = \lambda\mathbf{x}$.

Eigenvalue. A value λ for the eigenvector \mathbf{x} above. Example: λ is a complex root value.

Vector space. Contains vectors with magnitude and direction, which can be added together and multiplied by scalars in a field.

Module. A module over a ring is a generalisation of a vector space over a field, being an additive abelian group like a vector space, where the scalars are the elements of a ring.

14. Division algebra. (chapters III and V). A ring with division where multiplication might be nonassociative. Two elements of a division algebra cannot be multiplied giving zero unless one of them is zero.

Quaternions. (chapters III and V). A type of associative division algebra.

Exquaternions. (chapter V). A type of algebra obtained from the quaternions, neither associative nor with complete division.

Octonions, \mathbb{O} . (chapter V). A nonassociative division algebra.

Exoctonions. (chapter V) A type of algebra derived from the octonions, neither associative nor with complete division.

n-novanions. (chapter V). An n dimensional nonassociative division algebra, but not when both the real parts in a multiplication are zero.

15. Norm. Applied to complex numbers $a + bi$, the norm is $\sqrt{a^2 + b^2}$. For intricate numbers $a1 + bi + c\alpha + d\phi$ the norm squared is $a^2 + b^2 - c^2 - d^2$. Applied to a $n \times n$ matrix B , the norm is the positive n th root of $\det B$. Applied to n-novanions $a1 + bi + c\alpha + d\phi + b'i' + c'\alpha' + d'\phi' + \dots$, the norm is $\sqrt{(a^2 + b^2 + c^2 + d^2 + b'^2 + c'^2 + d'^2 + \dots)}$.

Interlayer operator \underline{v}_p . (chapter IV).

Diamond operator \diamond . (chapter IV).

Left roll operator s° . (chapter IV).

Right roll operator $^\circ s$. (chapter IV).

Split product. (chapter IV).

16. Standard protocol. (chapter VII). The ordinal infinity $\Omega_N = \sum_{\text{all } N} 1$. This is not a natural number, and is treated as being irreducible.

Ladder number. A superexponential expression in Ω_N , with Eudoxus coefficients.

Strict transfer principle. The axioms for variables in a superexponential algebra also hold for the variable Ω_N .

Winding number. The number of times a loop winds round a point.

17. Additive format of a polynomial equation. The form $ax^n + bx^{n-1} + \dots + d = 0$. (chapters VII and VIII).

Monic polynomial. Example in the case of a polynomial equation: when a above $= 1$.

Fundamental theorem of algebra. The complex polynomial in additive format given by $ax^n + bx^{n-1} + \dots + d$ always has some values which are zero.

Multiplicative format of a polynomial equation. The form $(x - p)(x - q) \dots (x - t) = 0$.

Zero of a polynomial. A value of a polynomial $f(x) = ax^n + bx^{n-1} + \dots + d$ so that $f(x) = 0$.

Root of a polynomial equation. The roots of a polynomial equation $f(x) = 0$ are the values of x satisfying this.

Degree of a polynomial. The value of n for $f(x)$.

Duplicate root. A root of the equation $(x + a)^2 = 0$.

Antiduplicate root. A root of the equation $(x + a)(x - a) = 0$.

Independent roots. Occur when no known dependency relation is used in the solution of a polynomial equation.

Dependent roots. Occur when a known dependency relation is used in the solution of a polynomial equation.

Polynomial entity. A polynomial equation with dependent roots.

Multivariate polynomial. A polynomial in a number of variables.

Variety. A polynomial equation in a number of variables. Example: $3x^2y + xyz + 4x^2z^2 = 0$.

18. Equivalence relation \equiv in a set S . Satisfies $a \equiv a$ (*reflexive*), if $a \equiv b$ then $b \equiv a$ (*symmetric*) and if $a \equiv b$ and $b \equiv c$ then $a \equiv c$ (*transitive*), for $a, b, c \in S$.

Equivalence class. A partition of a set where an equivalence relation between elements defines membership of the partition.

Partial order \leq of a set S . Satisfies $a \leq a$, if $a \leq b$ and $b \leq a$ then $a = b$ (*antisymmetric*) and if $a \leq b$ and $b \leq c$ then $a \leq c$, for $a, b, c \in S$.

Total order \leq of a set S is a partial order existing for all $a, b, c \in S$.

Well-ordering \leq of a set S . A total order where every nonempty subset has a least element.

19. Left (or right) coset of a subgroup S of G is the set of elements aS (or respectively Sa), with $s \in S$ and $a \in G$.

Quotient group G/S of $G \bmod S$. The family of left cosets of the group G with subgroup S , sG , $s \in S$.

20. Ideal, C . (chapters III and XI). A subset of a ring, A . $\{c, d\} \in C$ and $a \in A$ implies that $(c - d) \in C$ and both ac and $ca \in C$.

Principal ideal, (a) . The ideal generated by one element, a , of the ring A . For every $r \in A$, (a) is ra . Example: for $a \neq 0$ belonging to the integers \mathbb{Z} , $(3a) \subset (a) \subset \mathbb{Z}$.

Prime ideal, P . If a and b are two elements of A such that their product ab is an element of P , then a or b is in P , and P is not equal to the whole ring A . Example: integers containing all the multiples of a given prime number, together with zero. Example: the zero ideal (0) .

Maximal ideal, M . In any ring A , this is an ideal M contained in just two ideals of A , M itself and the entire ring A . Every maximal ideal is prime. Nonexistence: the zero ideal (0) is not a maximal ideal of \mathbb{Z} because $(0) \subset (2) \subset \mathbb{Z}$, nor is the ideal (6) , since $(6) \subset (2) \subset \mathbb{Z}$.

Nilradical, $N(A)$. The intersection of all prime ideals of a ring.

Jacobson radical, $J(A)$. The intersection of all maximal ideals of a ring.

21. Monomial (chapter XI) in a variety is a term without the coefficient. Example: x^4y^2z .

Monomial order of monomials in a variety. For example: $1 < x^4y^2z$ and if $x^4y^2z < x^4y^3z^2$, then $(x^4y^2z)(x^ay^bz^c) < (x^4y^3z^2)(x^ay^bz^c)$.

Lexicographic order $<_{\text{lex}}$ of monomials in a variety. The monomial order where the first powers g, h that differ satisfy $g < h$. Example: $x^4y^gz < x^4y^hz^2$, with $g < h$.

Degree lexicographic order $<_{\text{deglex}}$ of monomials in a variety. For example: $x^ay^bz^c <_{\text{deglex}} x^dye^zf$ if $a + b + c < d + e + f$, and if they are equal, then revert to lexicographic order.

Degree reverse lexicographic order $<_{\text{degrevlex}}$ of monomials in a variety is defined by $x^ay^bz^c <_{\text{degrevlex}} x^dye^zf$ if $a + b + c < d + e + f$, and if they are equal, then revert to right to left lexicographic order with the degrees interchanged. Example: $x^2y^2z^3 <_{\text{degrevlex}} x^4yz^2$ so the total degrees are equal, but then order with 3 and 2 in z interchanged.

Gröbner basis. A set of multivariate polynomial divisors of a multivariate polynomial with unique remainder.

22. Open set. Example: the interval $a < x < b$ with the end points a and b removed.

Closed set. Example: the interval $a \leq x \leq b$ with the end points a and b present.

Topology. A theory of space using open and closed sets.

Homotopy. A theory of paths through a topological space.

23. Exact sequence. (chapter III).

Homology. A theory of holes. The dimension of the n th homology is the number of holes in a space for dimension n .

24. Explanation. A theory or theorem using matrices or other combinatorial means.

Charade. The image of an explanation in homological algebra.

Deconstruction. The mathematical refutation of a generally accepted result.