

CHAPTER I

The meaning of the finite and the infinite

1.1. Introduction.

We give a description of formal language, and will use it to express the axioms of modified Zermelo set theory with the axiom of choice (mZFC). These axioms have been given in [Ad15] volume I, chapter III, section 2, and [Ad15] Volume II, chapter XIV, sections 2 and 5. We then extend the Peano axioms which define the rules for standard arithmetic.

Transfinite number theory is considered for the transnatural numbers \mathbb{M}_t , where the theory is an extension of finite arithmetic, and this is then used to define the real numbers in a similar way. We repeat a proof given in [Ad15] on the inconsistency of the uncountable continuum hypothesis with respect to the countability of the rationals. Then we extend results on the ladder algebra of ordinal arithmetic given in [Ad15] to \mathbb{M}_t .

The definition of real numbers given here will be used in volume III, chapter VI, to give a second proof of the general Riemann hypothesis.

1.2. Formal language. [5Co66]

In 1908, Zermelo presented a formal set of axioms for set theory which covered all reasoning in mathematics. The axiomatisation of set theory was in keeping with the spirit of the school of Formalism, led by David Hilbert. In the Formalist point of view, mathematics was seen as a purely formal game played with marks on paper, where the game must not lead to an inconsistent result.

The symbols used in a formal language, which are not the minimal set of symbols in the interpretation system we will give, we will designate as follows

NOT	&	OR	\Rightarrow	\Leftrightarrow
not	and	or	implies	if and only if
\forall	\exists	=	(,)	x, '
for every	there exists	equals	parentheses	variable symbols

In principle, the names used under these symbols which give their meaning are independent of the formal language.

We have already met the symbols in the first row in chapter XII. They are connectives used in a logic called the *propositional calculus*. The symbols used in the combined first and second rows are the symbols of the *predicate calculus*. On the second row the symbol \forall , for every (or for all), is known as the universal quantifier, and the symbol \exists , there exists (or for some), is the existential quantifier. Parentheses are used in the language to ensure its precise evaluation. When this evaluation is obvious, they may be omitted. The variable symbols will be formed as x, x', x'', etc., and we will often rewrite them as x_1, x_2, x_3 , etc. if that is useful.

From this set of symbols we can create others. Variable symbols with a special defined use in a system, such as the identity element for a group, are called constants, and will be denoted by c, c', etc. The variable symbols with the parenthesis symbol can be used to describe

relation symbols, R_1, R_2, R_3 , etc. Each R_k is provided with a natural number n_k which gives the number of variables in the parentheses which follow. To give examples of how these symbols could be used, for $n_0 = 2$ we can define a binary predicate $R_0(x_1, x_2)$, and this could represent $x_1 \leq x_2$, for $n_1 = 3$ we can define a ternary predicate $R_1(x_1, x_2, x_3)$, and this could represent $x_1 + x_2 = x_3$.

By these means symbols in the formal language can be used to express relations in arithmetic. For example if we wanted to express the uniqueness of addition in our language, this is

$$\forall x \forall y \forall z \forall u ((R_1(x, y, z) \& R_1(x, y, u)) \Rightarrow z = u).$$

Similarly, if $R_2(x, y, z)$ represents multiplication as $x \cdot y = z$, then we can express associativity of multiplication in the formal language by

$$\forall x \forall y \forall z \exists t \exists u \exists v ((R_2(x, y, t) \& R_2(t, z, u) \& R_2(y, z, v) \& R_2(x, v, u)).$$

Although the existence of the additive identity zero can be expressed as $\exists x \forall y (R_1(x, y, y))$, rather than restate this every time we use it, we will introduce the axiom $\forall x (x + 0 = x)$. We can now see a way of, say, describing the axioms of a field given in chapter III, section 4, by using formal language.

We now give the rules for writing grammatically correct statements in a formal language. These statements are called *well-formed formulas* (*wff's*).

- (1) Let x and y be variables, c and c' constants. Then $x = y$, $x = c$ and $c = c'$ are wff's.
- (2) If R is a relation symbol and u_1, \dots, u_n constants or variables, then $R(u_1, \dots, u_n)$ is a wff.
- (3) If A and B are wff's then so are NOT (A), (A) & (B), (A) OR (B), (A) \Rightarrow (B) and (A) \Leftrightarrow (B).
- (4) If A is a wff, so are $\exists x A$ and $\forall x A$.

The set of *subformulas* of a wff A is defined as the smallest set which contains A and is closed under the following rules

- (i) If NOT (B) is a subformula of A , so is B .
- (ii) If (B) & (C), (B) OR (C) or (B) \Rightarrow (C) are subformulas of A , so are B and C .
- (iii) If either $\exists x B$ or $\forall x B$ are subformulas of A , so is $B(c)$ for a constant c .

In subsequent developments we will need to distinguish between the ideas of *free* and *bound* variables. An example is as follows. Suppose we had $\forall x y = z$, which by rule (4) is a wff. Now $y = z$ does not involve x , so the quantifier $\forall x$ has no effect. A similar circumstance might be where variable and constant symbols occur in a wff without referencing anything.

Definition 1.2.1. A variable symbol in a wff is *free* or *bound* according as

- (a) Every variable occurring in a formula for rules (1) and (2) above is free.
- (b) Free and bound occurrences derived from rule (3) above are the same as those for A and B .
- (c) Free and bound occurrences of variables in $\exists x A$ and $\forall x A$ are the same as those for A except free occurrences of x are now called bound.

We allow binding to occur more than once. If this does not occur, the wff is called *good*.

Definition 1.2.2. A *statement* is a formula with no free variables.

Let us recall the rules for the propositional calculus given in chapter XIII of [Ad15]. We introduced two variables, τ , which we will now identify with true, and υ , which we identify with false.

Then the truth tables for statements A and B map $A(\tau, \upsilon)$, $B(\tau, \upsilon)$ to NOT A, A & B, A OR B, $A \Rightarrow B$ and $A \Leftrightarrow B$ as follows

NOT A	A & B	A OR B	$A \Rightarrow B$	$A \Leftrightarrow B$	A	B
υ	τ	τ	τ	τ	τ	τ
υ	υ	τ	υ	υ	τ	υ
τ	υ	τ	τ	υ	υ	τ
τ	υ	υ	τ	τ	υ	υ

The number of possible truth table functions is four for a function of one variable, 16 for a function of two variables, and in general 2^{2^m} for a function of m variables.

In fact, we can reduce the number of tables required to just NOT A and, say, A & B, since
 $A \text{ OR } B = \text{NOT}(\text{NOT } A \ \& \ \text{NOT } B)$
 $A \Rightarrow B = \text{NOT}(A \ \& \ \text{NOT } B)$
 $A \Leftrightarrow B = (A \Rightarrow B) \ \& \ (B \Rightarrow A).$

For the predicate calculus we can also introduce a simplification, since \forall can be replaced by NOT \exists NOT.

The objective of mathematics is to discover true theorems. The intention is that any valid statement is intuitively true in any interpretation of the relation and constant symbols used to form it. We use the term *valid* to describe statements formed by the rules we give next.

- (A) *Rule of the propositional calculus.* If P is a propositional function in the variables A_1, \dots, A_n that is always true, then on replacing any A_k by a valid statement results in a valid statement.
- (B) *Rule of inference.* This allows us to form new valid statements from old ones. If A and $(A) \Rightarrow (B)$ are valid statements, so is B.
- (C) *Rules of equality.* These allow us to manipulate equal signs. Let c, c' and c'' be constants.
 - (i) $c = c'$, $(c = c') \Rightarrow (c' = c)$ and $((c = c') \Rightarrow (c' = c'')) \Rightarrow (c = c'')$ are valid statements.
 - (ii) If A is a statement, and if A' represents A with every occurrence of c replaced by c', then $(c = c') \Rightarrow ((A) \Rightarrow (A'))$ is a valid statement.
- (D) *Change of variables.* If A is a statement and A' is obtained from A by replacing each occurrence of the symbol x by x', then $(A) \Leftrightarrow (A')$ is a valid statement. This also applies to good statements, as we have defined them.
- (E) *Rule of specialisation.* Let A(x) be a formula with one free variable x in which every occurrence of x is free, and A(c) be result from replacing every occurrence of x by the constant symbol c. Then $(\forall x A(x) \Rightarrow A(c))$ is a valid statement.
- (F) Let B represent a statement where c and x are absent. If $A(c) \Rightarrow B$ is valid, so is $\exists x A(x) \Rightarrow B$. This allows us to argue about a statement using an 'arbitrary' constant c, and infer that $\forall x A(x)$, since we have used no special properties of c.
- (G) *Rearrangement of quantifiers to the beginning of a statement.* Let A(x) have x as its only free variable, where every occurrence of x is free, and B be a statement with x absent.
 - (i) $(\text{NOT } (\forall x A(x))) \Leftrightarrow (\exists x \text{NOT } (A(x)))$.
 - (ii) $((\forall x A(x)) \ \& \ (B)) \Leftrightarrow (\forall x ((A(x)) \ \& \ (B)))$.
 - (iii) $((\exists x A(x)) \ \& \ (B)) \Leftrightarrow (\exists x (A(x)) \ \& \ (B))$.

1.3. Modified Zermelo-Fraenkel set theory (mZFC).

Sets are often used in the general description of mathematical systems; they include but are not restricted to collections of various types of number, for instance sequences or functions of numbers. The notion of a natural number itself can be expressed within this system. Further, we can introduce sets with an infinite number of members, or elements. We will not discuss in this section all aspects of generalisations of sets. Such systems can for example be defined in category theory, and some are known as toposes.

A set of axioms for set theory was given for the first time by E. Zermelo in 1908, which was later developed by A. Fraenkel. This combined theory is often called Zermelo-Fraenkel (ZF) set theory and can be used to describe all of traditional mathematics. When the axiom of choice is appended, it is known as ZFC. We list the axioms of mZFC, and comment on them. The symbol m indicates we have modified an axiom.

(1) *Axiom of extensionality.*

$$\forall x, y (\forall z (z \in x \Leftrightarrow z \in y) \Rightarrow x = y).$$

As for the system Z_2 , this states that a set is determined by its members.

We can now define the *includes* symbols.

$$x \subseteq y \Leftrightarrow \forall z (z \in x \Rightarrow z \in y) \text{ means } x \text{ is included in } y, \text{ and can equal it.}$$

$$x \subset y \Leftrightarrow x \subseteq y \ \& \ \text{NOT } x = y \text{ means } x \text{ is properly included in } y; \text{ it does not equal } y.$$

(2) *Axiom of the empty set.*

$$\exists x \forall y (\text{NOT } y \in x).$$

This empty set is denoted by \emptyset .

(m3) *Axiom of the void set.*

$$\exists x \forall y (\text{NOT } y \in x) \ \& \ y \in x).$$

This void set is denoted by \ominus , for which $\psi(x)$ defined later in (m7_n) is identically false.

(4) *Axiom of unordered pairs.*

$$\forall x, y \exists z \forall w (w \in z \Leftrightarrow w = x \text{ OR } w = y).$$

The set z of unordered pairs is denoted by $\{x, y\}$. This means that $\{x\}$ is $\{x, x\}$. For an ordered pair or *Cartesian product* $\langle x, y \rangle$, we define this as the set $\{\{x\}, \{x, y\}\}$. It is possible to prove that $\langle x, y \rangle = \langle u, v \rangle$ implies $x = u$ and $y = v$.

A definition of a *function* can now be given.

A function is a set f of ordered pairs such that $\langle x, y \rangle$ and $\langle x, z \rangle$ in $f \Rightarrow y = z$. The set of x is called the *domain* of the function and the set y the *codomain, image* or *target*. Some authors define the range to be the codomain of a function and others the domain, so we do not use this. A *map* or *mapping* is the set corresponding to the function itself. This mapping is *one-to-one*, an *injection* or a *monomorphism* (the last of such triples are used particularly for groups) if every $x' \in x$ maps to a unique $y' \in y$. A map is *onto*, a *surjection* or an *epimorphism* if every $y' \in y$ is the codomain of the function f with domain x . The mapping is *one-to-one and onto*, a *bijection* or an *isomorphism* if it is both injective and surjective.

(5) *Axiom of the union of sets.*

$$\forall x \exists y \forall z (z \in y \Leftrightarrow \exists u (z \in u \ \& \ u \in x)).$$

We call the set y the union of all the sets in x . Using axiom (4) we can then show that given x and y , there exists a z satisfying $z = x \cup y$, that is, $u \in z \Leftrightarrow u \in x \text{ OR } u \in y$.

(6) Axiom of *infinity*.

$$\exists x (\emptyset \in x \ \& \ \forall y (y \in x \Rightarrow y \cup \{y\} \in x).$$

We have already come across this in Z_2 : if x is a natural number, the successor of x is defined as $x \cup \{x\}$.

(m7_n) Axiom of *extended comprehension* (also called the modified axiom of *replacement*).

$$\forall u_1, \dots, u_k (\forall x \exists! y A_n(x, y, u_1, \dots, u_k) \Rightarrow \forall v \exists w \forall z (z \in w \Leftrightarrow \exists r ((r \in v \ \& \ A_n(r, z, u_1, \dots, u_k)) \text{ OR } r \in \odot))).$$

This axiom scheme says that if for fixed u_1, \dots, u_k $A_n(x, y, u_1, \dots, u_k)$ defines y uniquely as a function $\psi(x)$ of x , then for each v the codomain of v mapped to $\psi(v)$ creates a set, which may be void. Whenever we define a set by “ S is a set such that ...” we are using this axiom. Here the property A_n which defines ψ may be very non-constructive or inconsistent, for instance to verify it we may need to answer a question about infinite sets.

(8) Axiom of the *power set*.

$$\forall x \exists y \forall z (z \in y \Leftrightarrow z \subseteq x).$$

This states that for every x there exists a set y of all subsets of x . This y is defined by a property which is not covered by the axiom of restricted comprehension, because it is not defined as the codomain of any function.

(m9) Axiom of *consistent choice*.

If $n \Rightarrow A_n \neq \emptyset$ and $m \Rightarrow A_m \neq \emptyset$ are functions defined for all $n, m \in x$, then there exist other functions $f(n), f(m)$, with $f(n) \in A_n, f(m) \in A_m$ & NOT $(n = m \Rightarrow f(n) = f(m))$.

This allows us to do an infinite amount of ‘choosing’ even without a property that would allow us to define the choice function which could otherwise be defined by axiom m7_n.

The modification we have introduced is that this choice must be consistent, or nonvoid.

(10) Axiom of *regularity* (or *well-founding*).

$$\forall x \exists y (x = \emptyset \text{ OR } (y \in x \ \& \ \forall z (z \in x \Rightarrow \text{NOT } z \in y))).$$

This axiom is somewhat artificial and we include it for technical reasons only. It is never used in conventional mathematics. It states that every set $x \neq \emptyset$ contains a minimal element with respect to the \in relation (but not \subseteq), because we want our sets built up from \emptyset where all descending chains with respect to \in terminate with \emptyset . This axiom is similar to the well-ordering property, except that \in is not an ordering because we could have $x \neq y$, NOT $x \in y$ and NOT $y \in x$.

We have defined sets by modified Zermelo-Fraenkel set theory with the axiom of choice, mZFC, where the *axiom of extended comprehension* holds: For every set B and every well-formed formula $\psi(x)$, either there exists an $x \in X$ if and only if $\psi(x)$ holds and $x \in B$, or the set B is the void set, \odot , for which $\psi(x)$ is identically false.

Thus if P is a logical property and relation, called a predicate, there exists a set $Y = \{x: Px\}$, where we allow $X = \odot$, the void set, which satisfies a false predicate. For example, if a set is defined as the set of all barbers in town, B , and $b \in B$ is defined by b cuts hair for everyone in town, and b also satisfies the property that b does not cut b ’s hair, then this is a contradiction and $B = \odot$. We do not take this as an antimony in the axiomatics of sets.

\odot can be expressed indirectly as the complement of \mathbb{V} . Nonvoid sets obey a predicate which is somewhere true. For instance, applied to the set $\{X: X \notin X\}$ (Russell’s paradox),

$$\odot = \{x: x \in \emptyset \text{ AND } x \notin \emptyset\}$$

satisfies the paradox. A universal set (or universe) is then the set for a true predicate, say

$$\mathbb{V} = \{x: x \in \mathbb{V} \text{ OR } x \notin \mathbb{V}\},$$

which is self-referential, in that \mathbb{V} defines \mathbb{V} , or as a further example

$$\mathbb{V} = \{x: x = x\}.$$

The empty set satisfies $\emptyset \notin \emptyset$. If a set is void, it is empty: $\odot \subseteq \emptyset$, since a void set satisfies more conditions than an empty one. But \emptyset satisfies the condition that no $x \in \emptyset$, and therefore $\odot \in \emptyset$ has a false $\psi(x)$, so $\odot = \emptyset$. Though we can form the set $\mathbb{V}' \{x: \text{every } x \in \mathbb{V}'\}$, it is both the set complement of \emptyset and \odot , so $\mathbb{V}' = \mathbb{V}$.

Thus mZFC allows the construction of the set of all sets without a contradiction arising from Russell's paradox, and we do not require a theory which makes the distinction between the set of all sets, which is normally called a *class*, and other types of set.

We will now give further comments on what happens if some of these axioms are dropped. If the axiom of extensionality does not hold then the system may contain atoms, which are sets x with $\forall y (\text{NOT } y \in x)$ but for which the sets x can be distinguished.

If the axiom of infinity does not hold, then we can take as a model of ZF the set of all finite sets which can be derived from \emptyset . Since the other axioms are also valid for finite sets, this axiom is independent of the other axioms of ZF, since axiom (5) becomes false for finite sets.

1.4. The Peano axioms for the transnatural numbers, \mathbb{M}_t .

The axioms for a set specify in the axiom of infinity that there exists a non-trivial set called the natural numbers. We will work with the set of natural numbers, $\mathbb{N}_{\neq 0} = \mathbb{N} = \{1, 2, 3, \dots\}$, $\mathbb{N}_{\cup 0} = \mathbb{N} \cup \{0\}$ and the integers \mathbb{Z} given by $\{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$.

Equality, $=$, satisfies the property that it is an *equivalence relation*, namely if m, n, p are members in a set then $=$ is

$$\text{reflexive: } m = m \tag{1}$$

$$\text{symmetric: if } m = n \text{ then } n = m \tag{2}$$

$$\text{transitive: if } m = n \text{ and } n = p \text{ then } m = p. \tag{3}$$

If it is not the case that $m = n$, then we write $m \neq n$.

If elements of a set S satisfy the property of being in an equivalence relation given by $=$, its application to elements which belong to S forms a *partition* of S :

if $m \in S$, then the intersection of all elements $n \in S$ which equal m , with the set of all $n' \in S$ which do not equal m is the empty set:

$$\text{for every } m \in S, \{n; n = m \in S\} \cap \{n'; n' \neq m \in S\} = \emptyset,$$

where also

$$\text{for every } m \in S, \{n; n = m \in S\} \cup \{n'; n' \neq m \in S\} = S.$$

The natural numbers satisfy the Peano axioms describing a recursive procedure to generate them.

$$1 \in \mathbb{N} \tag{4}$$

$$\text{for every } n \in \mathbb{N}, \text{ there exists an } s(n) \text{ interpreted as } (n + 1) \in \mathbb{N} \tag{5}$$

$$\text{there is no number } 0 \in \mathbb{N} \text{ with } s(0) = 1 \tag{6}$$

$$\text{for two numbers } m, n \in \mathbb{N}, s(n) = s(m) \text{ implies } n = m. \tag{7}$$

$$\text{(induction) a subset of } \mathbb{N} \text{ containing } 1 \text{ and } s(n) \text{ whenever } n \in \mathbb{N}, \text{ is } \mathbb{N}. \tag{8}$$

We can express in an axiom system for sets for any property P that is not self-referentially true, that for the set $\{x: P; x \in X\}$, there exists a set $\{y: \text{NOT } P; y \in Y\}$. Since by this means we can introduce uncountable sets, we can now establish uncountable induction.

Definition 1.4.1. The *transfinite natural numbers* \mathbb{M}_t , where t belongs to an index set which also satisfies the properties below, satisfy the following axioms

$$1 \in \mathbb{M}_t \quad (9)$$

$$\text{for every } m \in \mathbb{M}_t \text{ there exists an } s(m) \text{ interpreted as } (m + 1) \in \mathbb{M}_t \quad (10)$$

$$\text{there is no number } 0 \in \mathbb{M}_t \text{ with } s(0) = 1 \quad (11)$$

$$\text{for two numbers } m, n \in \mathbb{M}_t, s(m) = s(n) \text{ implies } m = n \quad (12)$$

$$\mathbb{M}_t \subset \mathbb{M}_{s(t)} \quad (13)$$

$$\mathbb{M}_t \text{ is not bijective to } \mathbb{M}_{s(t)} \quad (14)$$

$$\text{there is no proper subset } \mathbb{M}' \text{ with the above properties satisfying } \mathbb{M}_t \subset \mathbb{M}' \subset \mathbb{M}_{s(t)} \quad (15)$$

$$\text{(induction) a subset of } \mathbb{M}_t \text{ containing } 1 \text{ and } s(m) \text{ whenever } m \in \mathbb{M}_t \text{ is } \mathbb{M}_t. \quad (16)$$

Notation 1.4.2. \mathbb{M}_t with the number 0 appended to it is denoted by $\mathbb{M}_{t \cup 0}$.

Define $<$ by the property, if $m, n \in \mathbb{M}_t$, then $m < n$ if and only if there exists a $p \in \mathbb{M}_t$ so that $m + p = n$.

Definition 1.4.3. The *transrational numbers* $\mathbb{Q}_{\mathbb{M}_t}$, $t \neq 1$, are the set of numbers $\pm m/n$ where $m \in \mathbb{M}_{t \cup 0}$ and $n \in \mathbb{M}_t$.

Definition 1.4.4. A *transprime* number is a transnatural number which is only divisible without remainder by 1 and itself.

Definition 1.4.5. A *transinteger* is a transnatural number multiplied by 1, 0 or -1.

Definition 1.4.6. A *transcomposite* number is the product of either a transnatural number of transintegers or may be used in a more restricted sense as a product of a transnatural number of transnatural numbers.

Definition 1.4.7. The *transalgebraic* numbers \mathbb{A}_t are combinations of the non-imaginary parts of transnatural numbers satisfying the field and either the exponential or more generally superexponential axioms of chapter III, section 2.

1.5. A model for the real numbers.

The definition of a symbol as a universal object is given in chapter III, section 8, on category theory, definition 3.8.4.

Definition 1.5.1. The *real transnatural* numbers represented by a symbol \mathbb{S}_t , where t belongs to a non-imaginary index set, satisfy the axioms for \mathbb{M}_t except possibly axiom (15).

Definition 1.5.2. The *real transrational numbers* $\mathbb{Q}_{\mathbb{S}_t}$, $t \neq 1$, are the set of numbers $\pm m/n$ where $m \in \mathbb{S}_{t \cup 0}$ and $n \in \mathbb{S}_t$.

Definition 1.5.3. A *real transprime* number is a real transnatural number only divisible without remainder by 1 and itself.

Definition 1.5.4. The *real transalgebraic* numbers \mathbb{A}_{S_t} are combinations of the non-imaginary parts of real transnatural numbers satisfying the field and either the exponential or more generally superexponential axioms of chapter III, sections 4 and 5.

Now the jump in transcendence is not limited as t increases. Numbers in \mathbb{A}_{S_t} are then discrete in $\mathbb{A}_{S_{(t+1)}}$ and continuous in $\mathbb{A}_{S_{(t-1)}}$. The question arises whether there is a model where S_t exists, but there are numbers with no imaginary components which do not belong to \mathbb{A}_{S_t} for any t . In the general case the union of this set with \mathbb{A}_{S_t} we call the *real numbers*, and we will find in the next section that we can form explicit calculations with models of them, which are representable when no such t exists.

1.6. The capital Ξ function.

The capital Ξ function will be used in Volume III, chapter VI to give a second proof of the general Riemann hypothesis. To take an example, for the exponential operation we can form

$$a^1 = a,$$

but in a conventional real exponential algebra, for $a \neq 1$ there is no

$$1^b = a.$$

What we do is introduce a function $\Xi(a)$ so that

$$1^{\Xi(a)} = a.$$

This number has been introduced culturally in an analogous way to our introduction of division by multizeros in the zero algebras of *Superexponential algebra*, volume I, chapter III. The idea can be extended to superexponential operations.

An interesting feature of $\Xi(a)$ is that it gives an ordering with a value which is greater than all transnatural ordinals, and thus gives a model for transcendence, and therefore in a proper sense, for real numbers.

Another observation is that $\Xi(a)$ is directly computable with the algebra we already have to hand, since

$$1 = e^{i2n\pi},$$

and if we choose the classical complex algebra, this means

$$1^i = (e^{i2n\pi})^i = e^{-2n\pi}$$

and with

$$e^{\log_e a} = a,$$

this gives

$$a = 1^{\Xi(a)} = (e^{-2n\pi})^i e^{2\pi n \left(\frac{\log_e a}{2\pi n}\right)},$$

$$\Xi(a) = 1 + \frac{i}{2\pi n} \log_e a. \quad \square$$

1.7. The Euclidean algorithm.

Strictly speaking, before proving the division with remainder theorem, we need to prove that *any transnatural number n greater than 1 is either transprime or a product of transprimes*.

Proof. By the *method of induction*, if we wish to prove a statement for a positive transnatural number n , we can assume it has been proved for any number less than n . If $n = 2$, the theorem is proved. If n is transcomposite, it can be represented as ab , where both a and b are greater than 1 and less than n , but then we know by the induction method that a and b are either transprimes or the product of transprimes. \square


The *fundamental theorem of arithmetic* states that *this factorisation is unique up to ordering*.

Proof. If n is transprime, the theorem is proved. Suppose n is transcomposite and there are two factorisations

$$n = pqr \dots = p'q'r' \dots$$

Let the transprimes be ordered in increasing size. By an induction hypothesis, no transprime p, q or r, \dots can be the same as any of p', q', r', \dots otherwise we could divide n by it and get two representations of a smaller number, where we would continue the proof with this smaller number. Since n is transcomposite it consists of at least two transprimes, so $n \geq p^2$ and $n \geq p'^2$ which implies $n > pp'$ with strict inequality, since p and p' are unequal. Now form $n - pp'$. This has pp' as a factor, so dividing $pqr \dots$ by p, p' must be a factor of $qr \dots$, a contradiction. \square

Suppose we represent the integer parts of the transrational numbers by notches along a line, for instance



then any transrational number may be represented uniquely by a transinteger plus a transrational number q equal to or greater than 0 and less than 1. This is an example of the theorem given by Euclid, in book 7 of the Elements.

The algorithm may be restated as:

Every positive transnatural number n can be written uniquely in terms of a positive transnatural number w less than n multiplied by another transnatural number $k > 0$ with a unique remainder $0 \leq u < w$:


$$n = kw + u. \tag{1}$$

Proof. If w divides n , then $u = 0$ and we are done. Otherwise assume (1). If n comes between kw and $(k + 1)w$, then (1) holds with $0 < u < w$. \square

Using the theorem, we can develop an arithmetic for fixed w in which we only consider u above. This transfinite arithmetic is known as congruence, or clock arithmetic. The equation above is then written in the notation

$$u = n \pmod{w},$$

where mod stands for the modulus, introduced by Gauss in his *Disquisitiones Arithmeticae*, which can be depicted in the example diagram for $w = 5$:



1.8. Fermat's little theorem.

Theorem 1.8.1. Fermat's little theorem extended to the transnatural numbers states that for a transprime p , verifiable directly for $p = 2$

$$x^p - x = bp. \quad (1)$$

for some unique b dependent on x .

Proof. We prove this by induction. For $x = 0$

$$0^p - 0 = 0p.$$

Assume (1) holds. Then for $x \rightarrow x + 1$, by the binomial theorem and the transnatural primality of p , so p does not divide any denominator

$$(x + 1)^p - (x + 1) = x^p - x + px^{p-1} + [p(p-1)/2]x^{p-2} + \dots + 1^p - 1 = bp + cp$$

for some unique c . \square (2)

1.9. Euler's totient theorem.

Let u be any positive transinteger, and let the totient $\varphi(u)$ denote the number of positive transintegers, 1 included, which are coprime to u and not greater than u . Since primality makes sense for transnatural numbers, it also makes sense for relative primality.

By definition $\varphi(1) = 1$. Also if p is a transprime number

$$\varphi(p) = p - 1.$$

Next suppose u transcomposite, and let p, q, r, s, \dots be the different transprimes dividing u .

Consider the series of transintegers, $1, 2, 3, \dots, u$. Of these the following are multiples of p :

$p, 2p, 3p, \dots, (t/p) \cdot p,$
(u/p in all).

Write these down with the sign $+$. Similarly, write down all the multiples of q, r, s, \dots each with the sign $+$.

In the same series there are $u/(pq)$ multiples of pq . Write these down with the sign $-$, and do the same with all the multiples of pr, ps, qr, \dots (taking all the products of p, q, r, s, \dots two at a time). Next write down all the multiples of the triple products pqr, pqs, \dots each with the sign $+$, and so on, until at last we come to the multiples of $pqrs \dots$ with sign $(-1)^{k-1}$, k being the number of different primes.

Now take any number θ which is not greater than t and not coprime to it. It will involve in its composition a certain number (λ say) of the different transprimes p, q, r, \dots . How many times will it occur among the multiples already written down?

By the binomial theorem, the number of combinations of λ things taken v at a time is

$$\lambda!/[v!(\lambda - v)!].$$

Evidently, taking its appearances in the order of the sets of multiples, θ will occur for λ times, the binomial coefficient, for $v = 1$ with the sign $+$, then for $\lambda(\lambda - 1)/2$ times for $v = 2$ with the sign $-$, then $\lambda(\lambda - 1)(\lambda - 3)/3!$ times for $v = 3$ with the sign $+$, and so on.

If then we take the algebraic sum of all the sets, we have θ occurring with a coefficient

$$\lambda - \lambda(\lambda - 1)/2! + \lambda(\lambda - 1)(\lambda - 3)/3! - \dots = 1 - (1 - 1)^\lambda = 1.$$

Thus the algebraic sum in question is the sum of all positive transintegers not greater than u and not coprime to it. Now the *number* of these integers is equal to the excess of the number of positive terms in the whole sum, as originally written, above the number of negative terms:

$$u\{(1/p + 1/q + 1/r + \dots) - (1/(pq) + 1/(pr) + 1/(qr) + \dots) + (1/(pqr) + 1/(pqs) + \dots) - \dots + (-1)^{k-1} \cdot 1/(pqrs \dots)\}.$$

Subtracting this from u , we have finally

$$\varphi(u) = u(1 - 1/p)(1 - 1/q)(1 - 1/r) \dots \quad \square \tag{1}$$

Corollary 1.9.1. If u is odd, $\varphi(u)$ is even. \square

1.10. Transp-adic numbers.

The p -adic number system for any transprime number p extends the ordinary arithmetic of the rational numbers in a different way from the extension of the rational number system to the real and complex number systems. The extension is achieved by an alternative interpretation of the concept of *closeness* or *absolute value*. In particular, trans p -adic numbers have the property that they are considered closer the more their differences are divisible by higher powers of p . These numbers exist also for transprimes. A transp-adic number b is represented for $j \geq k \in \mathbb{Z}_t$ and $0 \leq a_i < p \in \mathbb{M}_t$ by

$$b = \sum_{i=j}^k \frac{a_i}{p^i}. \tag{1}$$

Congruence arithmetic can then be encoded if

$$(j - k) = 0 \pmod{q} \tag{2}$$

usually for some transprime q , but now an ordering is not naturally defined.

1.11. The inconsistency of the UCH property for $\mathbb{N}^{\mathbb{N}}$.

We repeat a small part of [Ad14] on countable and uncountable sets and ordinal infinities, and extend it, but first in this section we will find a rule for the conditions under which the principle of induction is valid for the properties of sets.

Doly García remarks that if a property holds for a set indexed by $1, n$ and $n + 1$, one cannot argue that it holds for the entire set, an instance being given by the set $A_1 = \{1\}$, with the property that $2 \notin A_1$, and in general for $A_n = \{1, 2, \dots, n\}$, where $n + 1 \notin A_n$, so that each complement exists for finite n , but not the entire set \mathbb{N} . This is part of the reason for us introducing ladder numbers, but we need to view this where nonstandard analysis is not used.

To respond to this criticism, if we look at the definition of the empty set just given, we notice that mZFC deals with predicates in a nonstandard way, in that the same sentence may range over true and false is admissible in the definition of a set, and false defines the void set in mZFC, and not otherwise. Then if we look at the complement of A_1 , this is $\mathbb{N} \setminus \{1\}$, and the complement of A_n is $\mathbb{N} \setminus \{1, 2, \dots, n\}$, so that applying the principle of induction, over the set $A_{\mathbb{N}}$ we are dealing with $\mathbb{N} \setminus \mathbb{N}$, which is the empty or void set, and thus the same property ranges over true and false in accordance with its principle of induction for sets.

However, as we will put in more technical detail after theorem 1.11.2

Proposition 1.11.1. *If the property for a set does not range over the void set, it does not become false.*

In mZFC, if a predicate holds for all finite $n \in \mathbb{N}$, either the predicate does not hold for all n , which could be the case for standard set theory, where the set of all n not including all finite n is empty, which satisfies a false predicate, or the predicate holds for all n , which is the case for nonstandard set theory, since now the set of all n not including all finite n has elements.

When mapping properties, for instance a bijection $\{\mathbb{V}, \emptyset\} \leftrightarrow \{\mathbb{V}, \emptyset\}$ restricted to $\mathbb{V} \rightarrow \mathbb{N}$, for standard set theory possible mappings are $\mathbb{V} \leftrightarrow \mathbb{V}$, $\mathbb{V} \leftrightarrow \emptyset$, $\emptyset \leftrightarrow \mathbb{V}$ and $\emptyset \leftrightarrow \emptyset$. Since \mathbb{V} maps to true and \emptyset to false, we may and will define that \leftrightarrow satisfies the truth table for IF and only IF, written as \Leftrightarrow , with $A \leftrightarrow B \leftrightarrow C$ satisfying $(A \Leftrightarrow B) \& (B \Leftrightarrow C)$.

$A \leftrightarrow B$	A	B
T	T	T
F	T	F
F	F	T
T	F	F

This allows us to state that the principle of induction applies to bijective properties in standard set theory. \square

If \mathbb{N} is countable, so is its power set $2^{\mathbb{N}}$, in general there is a bijection between \mathbb{M}_t and $2^{\mathbb{M}^t}$, \mathbb{S}_t and $2^{\mathbb{S}^t}$, and \mathbb{R}_t to $2^{\mathbb{R}^t}$, as an extension of the theorem proved here.

Theorem 1.11.2. *There exists a bijection between the set of rationals, \mathbb{Q} , and what we define as $\mathbb{N}^{\mathbb{N}}$.*

Proof. Nowhere does the induction we will carry out range over the predicate for a void set, since its function of functions $\mathbb{N} \rightarrow \mathbb{N} \rightarrow \dots$ mapped in sequence to $\mathbb{N} \rightarrow \mathbb{N} \times \mathbb{N} \rightarrow \dots$ has a fixed and not exhausted function as its domain. This assertion is proved in chapter XIV section 6 of [Ad15], where cardinals defined by bijections are shown to be lowest ordinals, ordinals are countable if there is a bijection to \mathbb{N} , and it is proved that because a union of countably many countable sets is countable, that superexponential operations on countable ordinals are countable.

For finite or countably infinite sets S , T or U we define \equiv by the existence of at least one bijection within the natural numbers \mathbb{N} . This is an equivalence relation, in that $S \equiv S$, if $S \equiv T$ then $T \equiv S$, and if $S \equiv T$ and $T \equiv U$, then $S \equiv U$, so this forms a partition between those sets belonging to the equivalence class, and those outside it. Then for sets S_n , T_n , $n \in \mathbb{N}$, if for each n $S_n \equiv T_n$, then for the set of all S_n , $\{S_{nn}\} \equiv \{T_{nn}\}$, where this means if $S_n \equiv T_n$, then the bijection is maintained for $S_{n+1} \equiv T_{n+1}$, and the second subscript indicates a distinguished copy for $S_n \neq \emptyset$, defined inductively: $S_{n1} = \{S_n\}$, $S_{n2} = \{\{S_n\}\}$, $S_{np+1} = \{S_{np}\}$. \square

In this work we will adopt the argument of Cantor that *the set \mathbb{Q} of rationals is countable.*

Define the Cartesian product of all natural number pairs $\mathbb{N} \times \mathbb{N}$ as \mathbb{N}^2 . Consider the rational numbers not in lowest terms given by the set $\mathbb{Q} \equiv \{\{1/n\}, \{2/n\}, \{3/n\}, \dots\} \equiv$ the unordered distinguished copies $\{\mathbb{N}_1, \mathbb{N}_2, \mathbb{N}_3, \dots\} \equiv \mathbb{N} \times \mathbb{N}$ (a set of ordered pairs) which by the Cantor argument given next is $\equiv \mathbb{N}$.

There is a bijection

$$\mathbb{N}_n \leftrightarrow \mathbb{N}$$

obtained by stripping out all containing parentheses. Thus

$$\mathbb{N}_n \equiv \mathbb{N}, \tag{3}$$

and we have proved the equivalence (2), where we can add the comment for any n ,

$$n \times \mathbb{N} \equiv \mathbb{N}.$$

From the equivalence (3) we get

$$\mathbb{N}_p^p \equiv \mathbb{N}. \tag{4}$$

Thus we have shown directly that equation (1) holds, where

$$\mathbb{N}^p \rightarrow \mathbb{N} \text{ is injective implies } \mathbb{N}^{p+1} \rightarrow \mathbb{N} \text{ is injective,}$$

so, because there is no void set in the mappings, by induction we obtain its consequence

$$\mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N} \text{ is injective,}$$

and since for the constant $\{1\}^{\mathbb{N}}$

$$\mathbb{N} \leftrightarrow \{1\}^{\mathbb{N}} \times \mathbb{N} \rightarrow \mathbb{N}^{\mathbb{N}} \text{ is injective}$$

we derive the result

$$\mathbb{N}^{\mathbb{N}} \equiv \mathbb{N}. \quad \square \tag{5}$$

Another way of stating this feature is that the uncountable continuum hypothesis is not an axiom that is independent of the countability of \mathbb{Q} and this rule of induction, and the countability of \mathbb{Q} naturally takes precedence over the continuum hypothesis.

Thus in a version of the second order logic developed here, all superexponential operations resulting in the construction of sets build countable sets from countable sets. This is a new conclusion, and must replace the uncountable continuum hypothesis. \square

To deal with objections to the above proof, it has been stated that the arguments we have given in [Ad14] are dependent on representable infinitesimals. But the proof we have given here is not dependent on representable infinitesimals, and we state, but do not yet prove, that if representable infinitesimals are inconsistent then so are the axioms for a field, since their existence is contingent on either the axioms for a field, or on the existence of infinitesimals themselves.

Secondly, it might be claimed that there is no technical basis for applying induction to sets. As we have seen in section 2 on formal language and the axioms for mZFC in section 3, this is not so. Further, the discussion in [Ad15], chapter XIV, connects set theory with arithmetic, covered in the Peano axioms of section 4 defining induction. This [Ad15] discussion extends these considerations to arithmetic with superexponential operations, with similar results.

The technical basis we have just discussed may be expressed in category theory, presented in chapter III of this volume. An idea is that of a symbol representing a universal object, which could be a set. But because exponentiation is not directly representable in category theory, arising from the fact that exponentiation is nonassociative, we have to use section 9 of that chapter also, which represents nonassociative operations in a standard invariant form.

Thirdly, the previous proof raises the issue of the status of the Cantor diagonal argument on the uncountability of the real numbers. Before dealing with this, we address its counterpart for finite sets, where the Cantor argument does not work. We extend this idea to infinite but countable sets.

The Cantor diagonal argument is deconstructed as follows. First note the example of a finite set consisting of elements ordered as (a, b) , where a and $b \in \{0, 1\}$, can be described as two finite sets E, F where $E = \{(0, 0), (0, 1)\}$ and $F = \{(1, 0), (1, 1)\}$. In this example we define *diagonal* E to be found from

$$\begin{matrix} \underline{(0, 0)} \\ (0, \underline{1}), \end{matrix}$$

that is, $(0, 1)$ as underlined, so $\text{NOT diagonal } E = \text{NOT } (0, 1) = (1, 0)$. So E is finite, and the fact that $\text{NOT diagonal } E \notin E$ (but $\in F$) does not show that $\{E, F\}$ is not a finite set. We have proved that *for a finite set, the diagonalisable elements form a subset of the finite set under an ordering.*

In the finite case the Cantor theorem holds that the power set (the set of subsets) of any set is larger than the set, but the *NOT diagonal* relation for finite sets does not give the correct semantic interpretation for infinity. The correct definition of a set S_m being finite is that it is empty or has all elements (s_1, s_2, \dots, s_m) with fixed $m \in \mathbb{N}$, and an infinite set is not finite. \square

Definition 1.11.3. A *Eudoxus number*, $u \in \mathbb{U}$, is a number bounded within $[m, n] \in \mathbb{N}_{\neq 0}$ for some $n > m$, so that there exists some other natural number m which when multiplied by u lies within $[m, n]$. By bounded, we mean it has an absolute value, possibly represented by a norm, and this absolute value is positive, even if u is not.

Proposition 1.11.4. A *Cauchy sequence*, which is normally held to represent real numbers representable at most by a convergent sum of a countably infinite set of rational numbers, satisfies the *Eudoxus property*.

Theorem 1.11.5. In the standard representation of real numbers as Cauchy sequences, the real numbers are countable.

Proof. For the diagonal argument for infinite sets, consider the list of Eudoxus numbers denoted in binary and indexed by $2n_i$

$$\begin{array}{c|c} 2\mathbb{N} & \text{The set of all diagonalisable Eudoxus numbers with respect to an ordering} \\ \hline 2n_i & u_i \end{array}$$

Firstly, we will consider a specific ordering of \mathbb{U} . By a similar argument to the finite case *diagonal* $\{u_i\} = \{\text{the Eudoxus number with the } i\text{th digit taken from } u_i\}$, so $\text{NOT diagonal } \{u_i\} \notin \{u_i\}$ is consistent, since $\mathbb{N} \equiv 2\mathbb{N} \equiv (2\mathbb{N} + 1) \equiv 2\mathbb{N} \cup (2\mathbb{N} + 1)$, and all Eudoxus numbers are indexed by $n_i \in \mathbb{N} \equiv \mathbb{N}_{\text{diagonal}} \cup \{n_i\}_{\text{nondiagonal}}$. In this ordering $\{n_i\}_{\text{nondiagonal}}$ is one element appearing after the diagonal terms, or can be inserted in first to show the bijection with \mathbb{N} . Thus the utility of the diagonal argument disappears under a specific ordering.

We will now consider a generic ordering covering all possible orderings of \mathbb{U} . Then the larger set of $\{n_i\}_{\text{nondiagonal}}$ now corresponds to the set of possible \mathbb{U} , so each instance of this diagonal can appear, say, as the first element of \mathbb{U} in a different ordering. Thus the utility of the diagonal argument now disappears under a generic ordering, and we are back to proving $\mathbb{N}^{\mathbb{N}} \equiv \mathbb{N}$, which we have already done, and so the Eudoxus numbers are countable. \square

This theorem is in conflict with the assertion that $\{0, 1\}^{\mathbb{N}}$ is uncountable, which in turn uses the unacceptable semantic definition that the *NOT diagonal* relation for countably infinite sets acts as a criterion for uncountability. However as mentioned by P. J. Cohen in [1Co63], [1Co64] ‘one can construct models in which the set of constructible reals is countable’. \square

1.12. Ordinal (ladder) arithmetic in \mathbb{M}_t .

Definition 1.12.1. We adopt the *standard protocol* for ladder algebra:

$$\Omega_{\mathbb{M}_t} = \sum_{\text{all } \mathbb{M}_t} 1. \quad (1)$$

For any $n \in \mathbb{M}_t$, $n < \Omega_{\mathbb{M}_t}$, so $\Omega_{\mathbb{M}_t} \notin \mathbb{M}_t$. We will treat $\Omega_{\mathbb{M}_t}$ as being irreducible. This means we do not split $\Omega_{\mathbb{M}_t}$ into noncontiguous components, or truncate or extend it. We adopt for $\Omega_{\mathbb{M}_t}$ the negation shown below of a property attributed by Archimedes to Eudoxus of Cnidus for finite natural numbers: the ordinal infinity $\Omega_{\mathbb{M}_t}$ is inaccessible with respect to n and obeys the rule

$$\text{for every } m \in \mathbb{M}_t \text{ and for } \Omega_{\mathbb{M}_t} \text{ there does not exist an } n \in \mathbb{M}_t: \Omega_{\mathbb{M}_t} < mn. \quad \square \quad (2)$$

Definition 1.12.2. We adopt the *strict transfer principle* for ladder algebra:

the axioms for a field or zero algebra hold with respectively $a\Omega_{\mathbb{M}_t}$, $b\Omega_{\mathbb{M}_t}$ and $c\Omega_{\mathbb{M}_t}$ replacing some or all of a , b and c in these axioms. \square

An example is $1\Omega_{\mathbb{M}_t} = \Omega_{\mathbb{M}_t}$.

Definition 1.12.3. Ladder transnatural numbers $\mathbf{L}_{\mathbb{M}_t\cup 0}$ are defined by

$$\mathbf{L}_{\mathbb{M}_t\cup 0} = \bigcup_m [\mathbb{M}_t\cup 0(\Omega_{\mathbb{M}_t})^m], m \in \mathbb{M}_t\cup 0.$$

The algorithmic proof of a proposition by induction: choose a start value, assume for n and then prove for $n + 1$, now extends to $n \in \mathbf{L}_{\mathbb{M}_t\cup 0}$ under the strict transfer principle. However, the Peano axiom of induction states that the natural numbers are unique, so we must augment it for $\mathbb{N}\cup 0$ by saying it contains no elements $(\Omega_{\mathbb{M}_t})^m$. In ladder algebra the complement of all \mathbb{M}_t is not empty, so an induction predicate can always be true, even in mZFC.

What is the status of finite proofs in this situation? We have seen by the axiom of the strict transfer principle above, that proofs by countable induction over \mathbb{N} reduce to finite proofs. Thus there exist some finite proofs over countably infinite sets.

We argue in terms of states and not processes. For infinite processes, obtained values may oscillate infinitely. There are at least two approaches that we can take to enable consistency.

The first looks at logical deduction. Valid reasoning based on the evaluation of finitely determined states by processes which terminate finitely are retained. Otherwise valid infinite reasoning is restricted, so that we may be able to obtain consistent results previously unavailable. We can find *preferred evaluations* of these infinitely determined states or those with infinite processes so that the induction procedure is restricted for these types. We adopt the following method employing arguments under the strict transfer principle. For reasoning using all \mathbb{M}_t , start with the first element of \mathbb{M}_t , which is 1, and employ *linear induction* by allocating either \mathbb{M}_t , or an ordered block of n elements which belong to \mathbb{M}_t , so that the same proof is valid in each block. *Nonlinear induction* can be defined for other partitions.

The second approach extends Boolean logic. General multivalued logics are discussed in volume II, chapter V. Thus, for example, for the infinite type of systems we have been considering, we could have multivalued logic operating on three states, ‘true’, ‘false’ and ‘oscillates’. A particular type of multivalued logic discussed in chapter XIII of [Ad15] is *probability logic*. A Boolean type logic with two states τ and υ can be extended so that its values are linear combinations of τ and υ , in particular retaining the boundary condition that both τ and υ exist within the logic. A linear probability logic contains the states $c\tau + (1 - c)\upsilon$

where c is, say, a real variable. It is a theorem that oscillating values may be allocated as the value $\frac{1}{2}\tau + \frac{1}{2}\nu$ in a linear probability logic.

The model used to interpret this logic is not here the statistical correlation approach given in chapter XIII of [Ad15], although its evaluation may be found as a limit process acting on its correlation with oscillating values.

In the ‘paradox of the liar’ the value of “ ‘A is valid’ and ‘A is invalid’ ” is false, and is not the same as “ ‘A is valid’ and ‘A is invalid’ is false ” (which is true) on keeping track of the level of nesting of the quotes and their implicit parentheses.

Since for us, logic also includes parentheses in its syntax, validity or invalidity of a formula may depend on the presence or absence of such parentheses, and this includes in recursive statements, which must have them.

X is invalid means X is invalid
 (X is invalid) is valid means X is invalid,

where similar and further nestings are possible. For example there is a possible chain

((X is invalid) is invalid) is invalid).

Thus there are an infinite number of states and their infinite countable evaluation

((X is invalid) is invalid) is invalid) ...

is equivalent to the evaluation

$$Y = \prod_{\text{all natural numbers } \mathbb{N}} -1.$$

For example, using the strict transfer principle to evaluate

$$Y = \prod_{\text{all } \mathbb{N}} -1, \tag{3}$$

partition \mathbb{N} into blocks of pairs starting from 1, then for each pair evaluate as a member of

$$W = (-1)_{\text{odd } \mathbb{N}} \times (-1)_{\text{even } \mathbb{N}},$$

and the product is always 1. Thus under the linear induction principle,

$$Y = 1,$$

and consequently the infinite countable evaluation equivalent to finding Y

((X is invalid) is invalid) is invalid) ...

evaluates to X which is valid. Its linear probability logic evaluation is $\frac{1}{2}\tau + \frac{1}{2}\nu$. \square

There exist ideals of these blocks, in which proofs of the above type live, for the example above $1.W$ and $-1.W$, and say cosets of the ideal $1.W$

$$0 + W, \text{ and } -2 + W,$$

where only the 0 coset is the preferred evaluation. It follows that the preferred evaluation of $\Omega_{\mathbb{M}_t}$ obtained additively is even and not odd, and its linear probability evaluates to

$$\frac{1}{2}(\text{even} = \tau)\tau + \frac{1}{2}(\text{odd} = \tau)\nu. \square$$

We have seen that an uncountable set \mathbb{R} is not generated recursively by any means from \mathbb{N} , even when it has a countably infinite set embedded within it. We say $\Omega_{\mathbb{R}}$ with uncountable \mathbb{R} is strictly less than ultrainfinity $n\mathcal{U}$, $n \in \mathbb{N}$, whereas there exist a hierarchy of ultrainfinities $n\mathcal{U}^m$, $m \in \mathbb{N}$, including superexponential types of this sort of expression, all of which are greater than $\Omega_{\mathbb{R}}$. This arises because $(\Omega_{\mathbb{R}})^{-1} = \epsilon_{\mathbb{R}}$ is not a multizero; no $a - a = \epsilon_{\mathbb{R}}$. \square

We can further introduce a sequence

$$\Omega_{\mathbb{N}} < \Omega_{\mathbb{R}_t} < \Omega_{\mathbb{R}_u} < \dots$$

in which there are no bijections of the type $\mathbb{R}_t \rightarrow \mathbb{R}_u$. It can now occur that, say, an algorithm halts in \mathbb{R}_u but not in \mathbb{R}_t , so \mathbb{R}_u is inaccessible with respect to \mathbb{R}_t . \square

The allocation of two values, True and False, to logical evaluation may be extended to an n-fold set of values. There is a relation and a difference between these values and recursive procedures.

In recursion, if we apply the NOT operator to say the property of being countable for a set, then we have seen we can substitute the ordinal infinity $\Omega_{\mathbb{M}_t}$ describing ladder numbers in the countably infinite set \mathbb{M}_t by the uncountable ordinal infinity $\Omega_{\mathbb{R}_t}$ of \mathbb{R}_t described by the standard protocol

$$\Omega_{\mathbb{R}_t} = \sum_{\text{all } \mathbb{R}_t} 1. \quad (4)$$

If we now apply recursively this procedure to $\Omega_{\mathbb{R}_t}$, we get a higher order uncountability of a set \mathbb{R}_u with ordinal infinity $\Omega_{\mathbb{R}_u}$. There is then a countable or uncountable sequence

$$\Omega_{\mathbb{N}} \rightarrow \Omega_{\mathbb{R}_t} \rightarrow \Omega_{\mathbb{R}_u} \rightarrow \dots$$

corresponding to sets with successively higher cardinality.

$$\mathbb{N} \rightarrow \mathbb{R}_t \rightarrow \mathbb{R}_u \rightarrow \dots$$

For the allocation of NOT to the value True or False, we merely switch their values. Nevertheless, we can consider this mapping as a recursion (mod 2). For an n-valued logic this may be considered as recursive (mod n).

This comparison we have been making has up to now been restricted to algebras which are abelian and associative. It is our thesis that these ideas can be extended to nonabelian and nonassociative n-fold logics. \square

The axioms for exponentiation are given as a case of those for superexponentiation in chapter XVII of [Ad15], and in chapter II of this volume. We extend the strict transfer principle to these axioms too. It follows that we may employ $(\Omega_{\mathbb{M}_t})^2$ and $(\Omega_{\mathbb{M}_t})^v$ for v an exponent $\in \mathbb{F}$ or \mathbb{Y} . For v a positive number, $(\Omega_{\mathbb{M}_t})^v$ becomes an example of a hyperinfinity.

We will denote $(\Omega_{\mathbb{M}_t})^{-1}$ by $\epsilon_{\mathbb{M}_t}$. It follows from the property of $\Omega_{\mathbb{M}_t}$ that the ordinal infinitesimal $\epsilon_{\mathbb{M}_t}$ satisfies

$$\text{for every } m \in \mathbb{M}_t \text{ and for } \epsilon_{\mathbb{M}_t} \text{ there does not exist an } n \in \mathbb{M}_t: n\epsilon_{\mathbb{M}_t} > m. \quad (5)$$

Similarly $(\Omega_{\mathbb{M}_t})^{-u} = (\epsilon_{\mathbb{M}_t})^u$ is a hyperinfinitesimal. \square

If we define $\Omega_{\mathbb{Z}_t} = \sum_{\text{all } \mathbb{Z}_t} 1$, then

$$\Omega_{\mathbb{Z}_t} = 1 + 2\Omega_{\mathbb{M}_t}. \quad \square \quad (6)$$

The set \mathbb{Q} of zero or positive rational numbers where $q = m/n \in \mathbb{Q}$, $m \in \{\mathbb{M}_t \cup 0\}$, $n \in \mathbb{M}_t$, ignoring the equivalence relation where m/n is in lowest terms, may be mapped bijectively to the set of pairs $\{m, n\}$. Then under the axiom system we are adopting, derived from the strict transfer principle acting on all pairs bounded by m and n, the number of elements of \mathbb{Q} in not necessarily lowest terms is $\Omega_{\mathbb{M}_t} (\Omega_{\mathbb{M}_t} + 1)$. \square

Using the strict transfer principle we obtain the Euler formula

$$e^{i\theta\Omega} = \cos(\theta\Omega) + i\sin(\theta\Omega). \quad \square$$

We may obtain integrals using this principle:

$$\int_0^\Omega x^2 dx = \frac{1}{3} \Omega^3. \quad \square$$